

# NOT ALL ST ELEVATIONS ARE INFARCTIONS: MYOCARDIAL BRIDGING IN DEXTROCARDIA

Loveneet Kaur, MD; VonPatrick Delarosa, MD; Matthew Frank, MD

## Introduction

Myocardial bridging (MB)—most commonly involving the mid-left anterior descending artery (LAD)—can cause dynamic systolic compression, impaired diastolic filling, and demand–supply mismatch that clinically mimics acute coronary syndrome (ACS). Vasospasm and tachycardia may exacerbate ischemia in susceptible patients. Current reviews recommend beta-blockers or non-dihydropyridine calcium channel blockers as first-line therapy, with revascularization reserved for refractory cases. Nitrates should be used cautiously as they may worsen systolic compression.

## Case Description

A 48-year-old man with possible untreated type 2 diabetes, dextrocardia, and asthma presented with continuous emesis followed by hours of non-radiating substernal “lump-like” chest pain and mild dyspnea. He was hemodynamically stable; initial high-sensitivity troponin was <6 ng/L. Electrocardiography showed anterior ST-segment elevation (V1–V2) and reciprocal ST depression in inferior leads, poor R wave progression, right axis deviation. He was transferred for emergent coronary angiography.

## Investigations

Left heart catheterization revealed a 40% mid-LAD occlusion representing myocardial bridge with dynamic systolic compression and no obstructive coronary disease. Chest pain resolved post-procedure, and serial troponins remained negative. Aspirin was continued, and prophylactic anticoagulation for venous thromboembolism was initiated. Further ischemia assessment using coronary CT angiography (CCTA) with or without CT-derived fractional flow reserve (CT-FFR) or functional stress testing to quantify ischemic burden and guide long-term management was recommended.

## Discussion

This case highlights a STEMI mimic due to MB without fixed coronary stenosis, likely potentiated by transient coronary spasm. Prior reports document MB-associated ST-elevation and even myocardial infarction despite angiographically normal coronaries, underscoring the contribution of vasomotor dysfunction. Dextrocardia complicates ECG interpretation; mirror-image right-sided precordial lead placement (V1–V6) is essential to obtain accurate tracings and avoid misclassification of anterior changes. Management of symptomatic MB after exclusion of ACS centers on heart-rate reduction with beta-blockers or non-dihydropyridine calcium channel blockers, modification of cardiovascular risk factors, and reserving invasive or surgical strategies (e.g., unroofing, stenting) for objectively demonstrated, refractory ischemia.

## Conclusion

In patients presenting with ST-elevation but angiographically normal coronaries, myocardial bridging with superimposed vasospasm should be considered. Resolution of symptoms with persistently negative biomarkers strengthens this diagnosis. Accurate ECG interpretation in dextrocardia requires correct lead placement, and downstream functional imaging (CCTA ± CT-FFR or stress testing) is critical for quantifying ischemia, guiding therapy, and avoiding unnecessary revascularization.

## References:

1. Myocardial Bridging: Diagnosis, Functional Assessment, and Management: *JACC* State-of-the-Art Review, Sternheim,D; Power, D; Samtani,R; et.al; *JACC*, Volume 78, Number 22, doi/ [10.1016/j.jacc.2021.09.859](https://doi.org/10.1016/j.jacc.2021.09.859)
2. Interplay Between Myocardial Bridging and Coronary Spasm in Patients With myocardial Ischemia and Non- Obstructive coronary Arteries: Pathogenic and Prognostic Implications; Montone, R; Gurgoglione, F; Rinaldi, R; et.al, *Journal of the American Heart Association*, doi/pdf/10.1161/JAHA.120.020535
3. The myocardial bridge: incidence, diagnosis, and prognosis of a pathology of uncertain clinical significance; Santucci, A; Jacoangeli, F; Cavallini, *European Heart Journal Supplements*,2022 Nov 12;24(Suppl I):I61-I67. Doi [10.1093/eurheartjsupp/suac075](https://doi.org/10.1093/eurheartjsupp/suac075)
4. Comprehensive functional and anatomic assessment of myocardial bridging: Unlocking the Gordian Knot; Ciliberti, G; Laborante, Renzo; Francesco, M, et.al; *Front Cardiovasc Med*. 2022 Nov 8;9:970422. doi: [10.3389/fcvm.2022.970422](https://doi.org/10.3389/fcvm.2022.970422)
5. CT Fractional Flow Reserve for the Diagnosis of Myocardial Bridging-Related Ischemia: A Study Using Dynamic CT Myocardial Perfusion Imaging as a Reference Standard; Yu, Y; Yu, L; Day, X, Zhang, J; *Korean J Radiol*. 2021 Dec;22(12):1964-1973. <https://doi.org/10.3348/kjr.2021.0043>
6. Computed Tomography-Derived Fractional Flow Reserve: Developing A Gold Standard for Coronary Artery Disease Diagnostics; Hu,L; Wang, Y; Rao, J, et.al; *IMR Press*, doi [10.31083/j.rcm2510372/htm](https://doi.org/10.31083/j.rcm2510372/htm)
7. ECG Diagnosis: Dextrocardia; Mozyan, C; Levis, J; *Perm J*. 2019 Aug 15;23:18-244. doi: [10.7812/TPP/18.244](https://doi.org/10.7812/TPP/18.244)
8. Echocardiographic changes and quality of life after surgical unroofing of myocardial bridges; Bianco, F; Bucciarelli, V; Surace, F, et.al; *J Int Med Res*. 2021 May 20;49(5):03000605211014847. doi: [10.1177/03000605211014847](https://doi.org/10.1177/03000605211014847)